Grip Strength: The Essential Biomarker for Longevity

Imagine if a simple handshake could reveal vital insights into someone's healthspan and overall vitality.

Grip strength has emerged as a powerful biomarker for biological age, surpassing traditional metrics such as blood pressure in predicting risk of chronic disease, health outcomes and longevity. Its value extends beyond being a measure of frailty or physical performance, reflecting key health metrics - muscle mass, bone density, and balance - that are critical for maintaining functional independence.

Thus, grip strength is not only a measure of physical performance but also a key indicator of one's biological resilience, with significant implications for long-term well-being.

As healthcare systems shift towards proactive, preventative healthcare and scientific research focusing on defining more accurate indicators of biological and functional age, the simplicity and predictive power of grip strength testing make a compelling case for its inclusion in routine health assessments.

This document outlines how grip strength measurements enable effective health monitoring, guide lifestyle modifications and proactive interventions to optimize physical and cognitive function, delay age-related diseases, and enhance long-term quality of life.

"Not enough can be said about the importance of grip strength as you age. It's one of the strongest physical associations with longer life."

Peter Attia MD, Longevity Expert



Selective Highlights

1. Biological Aging

Marker of Biological Aging: Grip strength provides a measure of biological age, with **lower** grip strength correlating with faster DNA methylation age acceleration, a key marker of cellular aging [Peterson et al., 2022].

Indicator of Genetic Variability: Variations in grip strength may reflect **genomic differences in myotube differentiation and muscle repair,** offering insights into personalized aging trajectories [Matteini et al., 2016].

Inflammation and Aging ("Inflammaging"): Chronic inflammation, often associated with aging, accelerates muscle weakness, biological aging, and disability. **Grip strength serves as a key indicator of inflammation-related risks**, linking low strength to higher rates of multimorbidity and age-related decline [Strasser et al., 2018].



Early Mortality Predictor: Grip strength is a stronger predictor of mortality and cardiovascular events than systolic blood pressure, with weaker grip strength linked to a higher risk of early death across all income levels [Leong et al., 2015].

Cancer Mortality in Women: Women aged 40–69 with grip strength below 24.3 lbs* face a 61% higher risk of breast cancer-related mortality, highlighting its value in assessing cancer prognosis [Celis-Morales et al., 2018].

Lifespan Insights in Men: Men with the weakest grip strength in their late teens are 20% more likely to die by their mid-50s compared to those with moderate or high grip strength [Ortega et al., 2012].



3. Functional Fitness & Frailty

Early Indicator of Frailty and Sarcopenia: Grip strength below 44.1 lbs* for men and 28.8 lbs* for women is a key marker of frailty under the Fried Frailty Phenotype (FFP), while levels below 41.0 lbs* for men and 24.3 lbs* for women are diagnostic criteria for sarcopenia, according to the The European Working Group on Sarcopenia in Older People [Fried et al., 2011; Cruz-Jentoft et al., 2019].

Postmenopausal Screening for Osteoporosis: Weak grip strength is associated with reduced bone density and a higher likelihood of falls and fractures, particularly in post-menopausal women, making it an effective tool for early osteoporosis and fall risk assessment. [Li et al., 2018]

Preserving Independence Through Early Detection: Grip strength testing identifies functional decline caused by advancing muscle atrophy, which contributes to disability and diminished quality of life. Early interventions, including resistance training or muscle-strengthening activities for 30–60 minutes weekly, reduces the risk of all-cause mortality, cardiovascular disease, and cancer by 10–17% [Momma et al., 2022] and can improve strength, maintain independence, and combat sarcopenia [Strasser et al., 2018].



4. Chronic Disease

Cognitive Health and Dementia Risk: Lower grip strength is linked to a nearly two-fold increased risk of cognitive decline and dementia (HR=1.99; Cui et al., 2021). Higher strength reflects neuromuscular integrity and resilience to oxidative stress, potentially protecting cognitive function [Weaver et al., 2002].

Cardiovascular & Respiratory Disease: Grip strength serves as a marker for risk of cardiovascular disease, type 2 diabetes, hypertension, and multimorbidity. **Including grip strength in risk** scores improves accuracy in predicting mortality and heart disease, while a grip strength of 62.4 lbs* reduces respiratory disease mortality risk by 51% [Cheung et al., 2013; Celis-Morales et al., 2018; Mey et al., 2022].

*Note that all cut off points have been converted to GripAble-equivalent measurements for consistency.

Grip Strength In Practice

Clinical Application

- Use grip strength as a simple screening tool during routine check-ups, to track trends and assess risk factors. Prioritize early detection and treatment of sarcopenia to prevent adverse health outcomes, as recommended by The European Working Group on Sarcopenia in Older People.
- For high-risk patients, consider providing a hand dynamometer for regular, at-home monitoring.
- Pay attention when grip strength drops below appropriate cut-off points.

Patient Empowerment

- Share grip strength scores with patients as a clear, relatable measure of functional independence that they can understand and actively improve with guidance.
- Early detection of low grip strength enables targeted interventions, such as exercise programs and nutritional support, to enhance healthspan.

Practical Considerations

- Incorporate regular grip strength measurements alongside standard health assessments to provide objective data on age-related decline and healthspan.
- Use grip strength as a safer, more practical alternative to gait speed tests for assessing falls risk, especially when gait speed testing may compromise patient safety.
- Use grip strength to complement patient-reported outcomes, delivering actionable insights for both clinicians and patients.

Impact

Personalized & Optimized Healthspan

Grip strength serves as a holistic maker of vitality, enabling clinics to create personalized programs that enhance physical and cognitive performance.

Improved QALYs & Functional Independence

Grip strength assessments offer early detection of age-related decline, allowing interventions that preserve functional independence and extend quality-adjusted life years (QALYs).

Reduced Healthcare Burden

Grip strength assessments enable early identification and interventions for at-risk patients, thus mitigating the burden of managing chronic conditions reducing hospitalizations and long-term care needs.



Able Assess

The Only Grip Strength Platform

Equipped for **Data-Driven Healthcare**.

Designed to unlock the full potential of grip strength as a vital health biomarker, Able Assess meets the demand for high-quality, data-driven healthcare.

Able Assess is a groundbreaking, digitally enabled platform for grip strength measurement.

It combines our advanced digital dynamometer sensor technology with a comprehensive integrative platform and the most up-to-date normative dataset on the market to empower clinicians with actionable information that traditional tools simply can't offer.

What's Next?



Explore: Dive into our extensive library of studies on grip strength and longevity [here].

<u>0</u> =	FI V
⊙=	Ŵ

Understand: Get the essentials on hand dynamometry and how to integrate grip strength into your practice with our comprehensive guide.



Contribute: Partner with us in research or patient case studies to advance the knowledge of grip strength as a biomarker.



Get in touch: Website: www.able-care.co Email: hello@able-care.co